

PREDIABETES and DIABETES

Risk Screening Tool

Patient Name: _____ DOB: _____ Age: _____
 Smoker? Yes No (circle one) Gender: M F Ht: _____ Wt: _____ BMI: _____

Date of Visit:									
ADULTS TO SCREEN	All adults > 45 years of age. Check plasma glucose. Especially if BMI ≥ 25 ♦ If Normal and pt. has no risk factors (listed below) <u>Rescreen every 3 years</u>								
	Adults < 45 years of age. Check Plasma glucose: ♦ If BMI ≥ 25 and with one additional risk factor *								
	If results are abnormal, obtain 2nd test on a different day to verify diagnosis of Pre-DM or DM. <u>Rescreen every 1-2 years</u>								
*RISK FACTORS	Habitually inactive								
	First degree relative with diabetes								
	Member of a high risk ethnic population African American Native American Hispanic American Pacific Islander Asian American								
	Delivered a baby weighing >9 lbs or have been diagnosed with Gestational Diabetes								
	Hypertensive (≥140/90mmHg) or being treatment								
	Dyslipidemia (either) HDL < 35 mg/dl TG > 250 mg/dl								
	Has PCOS								
	Previously IFG or IGT								
	Other signs of insulin resistance (acanthosis nigricans)								
	Hx of vascular disease								
DIAGNOSTIC LAB TESTS	PRE-DIABETES								
	Impaired fasting plasma glucose (100-125 gm/dl on 2 different dates)								
	Impaired glucose tolerance (2 hrs after 75gm OGTT) (140-199 mg/dl on 2 different dates)								
	DIABETES								
PEDIATRIC	Fasting plasma glucose of ≥126mg/dl on 2 days OGTT 2 hour ≥200 on 2 days								
	Check fasting plasma glucose on at risk** youth >10 yoa, or at onset of puberty if puberty occurs at younger age:								
	**Overweight (BMI > 85th %tile for age and sex, or wt > 120% of ideal for ht. Plus 2 risk factors below)								
	**Family hx of type 2 diabetes in 1st or 2nd-degree relative								
	**High risk race or ethnicity (above)								
	**Signs of insulin resistance acanthosis nigricans Hypertension dyslipidemia PCOS								
<u>Rescreen every 2 years</u>									

Adapted from Kentucky Diabetes Network, 2006

Developed by Northern Michigan's Diabetes Outreach Network - TIPDON - 2007



Once Diagnosis of Pre-Diabetes is made, use Prediabetes Tracking Tool